

# Habitat differences in the food composition of the wasp-like spider *Argiope bruennichi* (Scop.) (Aranei : Araneidae) in Poland

Paweł Szymkowiak<sup>1</sup>, Piotr Tryjanowski<sup>2</sup>, Aleksander Winiecki<sup>3</sup>, Seweryn Grobelny<sup>3</sup> and Szymon Konwerski<sup>4</sup>

<sup>1</sup> Department of Animal Taxonomy and Ecology, Adam Mickiewicz University, Umultowska 89, 61-614 Poznań, Poland

<sup>2</sup> Department of Behavioural Ecology, Adam Mickiewicz University, Umultowska 89, 61-614 Poznań, Poland

<sup>3</sup> Department of Avian Biology and Ecology, Adam Mickiewicz University, Umultowska 89, 61-614 Poznań, Poland

<sup>4</sup> Department of Zoology, Agricultural University in Szczecin, Doktora Judyma 20, 71-466 Szczecin, Poland

Corresponding author : Paweł Szymkowiak, Department of Animal Taxonomy and Ecology, Adam Mickiewicz University, Umultowska 89, 61-614 Poznań, Poland; e-mail: szymkowi@amu.edu.pl

**ABSTRACT.** During the last few decades the wasp-like spider *Argiope bruennichi* Scopoli, 1772 expanded its wide geographical distribution in Europe. In this paper we describe and test differences in the diet composition of the wasp-like spider inhabiting a river valley (traditional habitat) and xerothermic grassland (new habitat) in Poland. From a total of 163 webs of *A. bruennichi*, 430 prey items were found, mainly insects : Coleoptera, Diptera, Homoptera, Heteroptera, Hymenoptera, Lepidoptera, Mecoptera, Odonata, Orthoptera and Neuroptera. Moreover, a semi digested specimen of the common frog *Rana temporaria* was recorded. Habitats differed significantly in the percentage distribution of eleven general food categories. Among potential influencing factors the number of caught prey was correlated only with the height of the web hub above the ground. The wasp-like spider is ecologically flexible in the use of novel food spectra and this probably allows the colonisation of new localities, as well as habitats.

**KEY WORDS :** *Argiope bruennichi*, diet, prey selection, web structure, frogs.

## INTRODUCTION

Amongst orb-weaving spiders the wasp-like spider *Argiope bruennichi* Scopoli, 1772 expanded its geographical range to northern parts of Europe. The range of the wasp-like spider was previously limited to south-east Europe (POETZSCH, 1963; BARABASZ & GÓRZ, 1998). However, during the last few decades the species has colonized many new sites in Belgium, France, The Netherlands, Poland, Denmark, Sweden and even Great Britain (GUTTMANN, 1979; JONSSON & WILANDER, 1999; MOYES, 1997; SCHARFF & LANGEMARK, 1997). The most important factors causing rapid geographical expansion of the wasp-like spider are : climate change, especially the increase in numbers of sunny and dry days in summer, floods of large rivers in Europe, as well as the establishment of large open habitats due to deforestation and drainage (DZIABASZEWSKI, 1959; GUTTMANN, 1979; HELSDINGEN, 1982; PUTS, 1989; WEICKMANN & GROBMEIER, 1997; SCHARFF & LANGEMARK, 1997; LINDEN, 2000).

Similar to other orb-weaving spiders, *A. bruennichi* is a generalist predator (RIECHERT & ŁUCZAK, 1982; RIECHERT & HARP, 1987; MALT et al., 1990; FASOLA, 1999). However, many studies showed that orthopterous insects and dragonflies were the main prey of the spider (URBAŃSKI, 1948; BEDNARZ, 1966; NYFFELER & BENZ, 1978; BARABASZ & GÓRZ, 1998). Moreover, CONRAD & BREINL (1992) even indicated a strong dependence between rich orthopteran fauna and distribution of *A. bruennichi* and linked the geographical range of the

wasp-like spider with orthopteran abundance, as potential food supply. On the other hand, some authors showed that other prey, such as dipterans, hemipterans (mainly aphids) and hymenopterans constituted an important part of the wasp-like spider diet (NYFFELER & BENZ, 1982; MALT et al., 1990). However, in publications on wasp-like spider diet only a simple description of food content is given, without linking it to habitat, potential food sources, or web structure.

Therefore, in this study we focused on (1) a description of the wasp-like spider's diet composition in two different habitats in Poland; (2) analyses of differences in prey caught in both habitats; (3) some factors affecting hunting success connected especially with web structure (e.g. stabilimenta). We discuss our findings in the light of colonisation of new areas (*sensu* localities, as well as habitats) in Poland (particularly) and in Europe (generally).

## MATERIAL AND METHODS

Studies were carried out during August 1999 in the Wielkopolska province, western Poland (52° N, 16° E). The study area covers two different habitats : wetland in the Warta river valley, near the village of Wrąbczykowskie Holendry (Poznań region) and small marginal grassland habitats (arable fields), near the villages : Powodowo, Obra, Daszewice (Wolsztyn and Poznań regions). The density of wasp-like spiders in both habitats is similar, ca 0.3 orb-webs/1m<sup>2</sup>.

The spider webs were monitored before noon, mainly 9-11 a.m. The geographical exposition was measured based on the axis of the web elevation to the ground level. To ensure that chosen prey items of *A. bruennichi* were sampled, only wrapped preys were removed from webs and stored in tubes with ethyl alcohol (70%). To explain silk details (stabilimenta and turns on stabilimenta) on webs, Fig. 1 was supplied. Because we did not control all parameters for all webs, sample size is different in various analyses. Standard statistical methods were used to describe and analyse the data (ZAR, 1999). All statistical tests were two-tailed. We considered  $P < 0.05$  as the minimum acceptable level of significance.

## RESULTS

### Prey structure

Sampling a total of 163 webs of *A. bruennichi* revealed 430 prey items (mean  $\pm$  SD =  $2.6 \pm 1.3$  prey/web). In the webs, insects from the following ten orders were represented: Coleoptera, Diptera, Homoptera, Heteroptera, Hymenoptera, Lepidoptera, Mecoptera, Odonata, Orthoptera and Neuroptera. The most numerous species caught in the webs were: Blue-tailed Damselfly *Ischnura elegans* (Van der Linden, 1832) and a squash bug *Coreus*

*marginatus* (Linnaeus, 1758). Moreover, in one web an eleventh category, one semidigested specimen of the common frog *Rana temporaria* (Linnaeus, 1758), was found.

Dragonflies and dipterans contributed over 62% of the total prey captures - Table I.




Fig. 1. – Web structure of *A. bruennichi*.

TABLE I

Composition of prey capture species in two different habitats.

Explanations: RV – river valley, XH – xerothermic habitats

Order	Family	Prey items	RV	XH	
Coleoptera	Chrysomelidae	<i>Agelastica alni</i> (Linnaeus, 1758)	1		
		<i>Cassida vittata</i> Villers, 1789	3		
		<i>Cassida viridis</i> Linnaeus, 1758	1		
		<i>Gastrophysa polygoni</i> (Linnaeus, 1758)		4	
		<i>Phyllotreta</i> sp.		2	
	Coccinellidae	<i>Coccinella septempunctata</i> Linnaeus, 1758	2		
		<i>Tytthaspis sedecimpunctata</i> (Linnaeus, 1761)		2	
		<i>Propylea quatuordecimpunctata</i> (Linnaeus, 1758)		1	
		Geotrupidae	<i>Anoplotrupes stercorosus</i> (Hartmann, 1791)	1	
		Cetoniidae	<i>Protœtia cuprea</i> (Fabricius, 1775)	1	
Diptera	Staphylinidae	Phalacridae		1	
		Latridiidae		8	
	Bibionidae		1		
		Chironomidae	2		
		Culicidae	3		
		Syrphidae	1		
		other	10	104	
		other	3	30	
Homoptera	Cercopidae				
Heteroptera	Coreidae	<i>Coreus marginatus</i> (Linnaeus, 1758)	20		
		Miridae	3		
		Nabidae	1		
	Pentatomidae	6			
	Scutellaridae	<i>Eurygaster</i> sp.	3		
	other			26	
Hymenoptera	Apoidea		7		
	Argidae		1		
	Formicidae		1		
	other		2	8	
Lepidoptera	Pyalidae	<i>Pleuroptya ruralis</i> (Scopoli, 1763)		6	
	other			11	
Mecoptera	Panorpidae	<i>Panorpa communis</i> Linnaeus, 1758	4		
Odonata	Coenagrionidae	<i>Ischnura elegans</i> (Van der Linden, 1823)	134		
	Libellulidae	<i>Sympetrum sanguineum</i> Müller, 1764	5		
	other			6	
Orthoptera	Tetrigidae	<i>Tetrix subulata</i> (Linnaeus, 1758)	1		
	Tettigoniidae	<i>Metrioptera roeselii</i> (Hagenbach, 1822)	1		
Neuroptera	Chrysopidae			1	
Anura	Ranidae	<i>Rana temporaria</i> (Linnaeus, 1758)	1		



1966; PFLETSCHINGER, 1976). Our findings (a small (0.5%) proportion of orthopteran prey items) suggest that especially in newly occupied habitats, the wasp-like spider could explore a new diet spectrum of Diptera and Hemipterans. NYFFELER & BENZ (1978), obtained similar results during a study of prey selection of the wasp-like spider on fallow land near Zurich (Switzerland). This suggests that the trapping of small prey is a possible adaptation to new habitats where much larger food items, e.g. grasshoppers and dragonflies are strongly limited. Although the wasp-like spider is behaviourally adapted (PASQUET & LEBORGNE, 1990, 1998) to catch orthopterans, the choice of smaller prey is an advantage to expand geographical distribution to the North (INDYKIEWICZ et al., 1995; BARABASZ & GÓRZ, 1998), as well as to occupy new habitats: small patches of xerothermic grassland in intensively used farmland, or even arable field (BARABASZ & GÓRZ, 1998; CICHOCKI, 1998; and our unpublished data).

Moreover, an additional adaptation to the occupation of a new habitat is a change of plant species used as substrate for silk attachment. For a congeneric species, *Argiope aurantia* Lucas, 1833, McREYNOLDS (2000) found that webs on grasses were more effective in trapping orthopteran prey than those established on herbs and composites. However, we did not test this relationship directly; we can only suggest that the use of a wide range of plants in new habitats improved possibilities of catching novel prey sources.

An important factor improving spider foraging strategy is web structure, especially conspicuous silk structures called stabilimenta (HERBERSTEIN et al., 2000). It is suggested that increased number of stabilimenta can attract prey to the web (STARKS, 2002). HERBERSTEIN (2000), in a study of *Argiope keyserlingi* Karsch 1878, showed that webs with more stabilimenta caught more prey than webs carrying fewer decorations. However, this author did not find a relation between web height above ground and the rate of prey capture. Our findings are in contrast to this. The number of prey caught by the wasp-like spider was correlated with web (hub) height above ground level. We found that higher webs were more successful. Furthermore, we did not find an effect of the number of stabilimenta on the number of prey items in the web. BLACKLEDGE & WENZEL (1999), studying dipterans as prey, obtained interesting results and found that this group of airborne insects were less frequently caught in webs with stabilimenta than in those without stabilimenta. Wasp-like spiders feed mainly on tiny invertebrates on the northern part of their distribution and our results show that Odonata and Orthoptera were caught very seldom. Therefore, this is probably the reason why stabilimenta are not needed to catch most of such spider prey. Selection pressure should cause a significant decrease in the percentage of webs with those ornaments and it should be tested during a long-term study in the future.

We found also the capture and partly ingestion of a small specimen of the common frog by the wasp-like spider. It is, according to our knowledge, the first record of foraging on a vertebrate by *A. bruennichi* (RIECHERT & LUCZAK, 1982; RIECHERT & HARP, 1987 and other cited literature). To fully understand the role of food in the col-

onisation of new localities and habitats this problem should be studied in detail, both by description of diet contents and by field experiments.

## ACKNOWLEDGEMENTS

We thank A. Gawroński for his help in data collection, S. Mielewczyk for his help in dragonfly identification, and V. Takacs, P. Zduniak and three anonymous reviewers for valuable comments on earlier versions of the manuscript.

## REFERENCES

- BARABASZ, B. & A. GÓRZ (1998). *Argiope bruennichi* (Scopoli, 1772) rare and insufficiently examined spider species in Poland. *Fragmenta Faunistica*, 41 : 255-267.
- BEDNARZ, S. (1966). New records of *Argiope bruennichi* Scop. (Argiopidae) in the Lower Silesia. *Przegląd Zoologiczny*, 10 : 179-185.
- BLACKLEDGE, T.A. & J.W. WENZEL (1999). Do stabilimenta in orb-webs attract prey or defend spiders? *Behavioral Ecology*, 10 : 372-376.
- CICHOCKI, J. (1998). The occurrence of *Argiope bruennichi* in crops. *Przegląd Przyrodniczy*, 9 : 113-114.
- CONRAD, R. & K. BREINL (1992). Beitrag zur Ausbreitung der Wespenspinne aus Ostthüringen. *Entomologische Nachrichten und Berichte*, 36 : 61-63.
- CROME, W. & I. Crome (1961). Paarung und Eiablage bei *Argiope bruennichi* (Scop.) auf Grund vor Freilandbeobachtung an zwei Populationen im Spreewald/Mark Brandenburg (Araneae : Araneidae). *Mitteilungen aus dem Zoologischen Museum in Berlin*, 37 : 189-250.
- DZIABASZEWSKI, A. (1959). The spider *Argiope bruennichi* Scop. in Poland in the light of new investigations. *Przyroda Polski Zachodniej*, 3 : 128-138.
- FASOLA, M. (1999). Experimental competition release in a community of web-weaving spiders. *Italian Journal of Zoology*, 66 : 153-158.
- GUTTMANN, R. (1979). Zur Arealentwicklung und Ökologie der Wespenspinne (*Argiope bruennichi*) in der Bundesrepublik Deutschland und den angrenzenden Ländern (Araneae). *Bonner zoologische Beiträge*, 30 : 454-486.
- HELSDINGEN van, P.J. (1982). Postglacial expansion of *Argiope bruennichi* Scop. now also as far as The Netherlands. *Levende Natuur*, 84 : 121-123.
- HERBERSTEIN, M.E., C.L. CRAIG, J.A. CODDINGTON & M.A. Elgar (2000). The functional significance of silk decorations of orb-web spiders: a critical review of the empirical evidence. *Biological Reviews*, 75 : 649-669.
- HERBERSTEIN, M.E. (2000). Foraging behaviour in Orb-Web Spiders (Araneidae) - do web decorations increase prey capture success in *Argiope keyserlingi* Karsch, 1878. *Australian Journal of Zoology*, 48 : 217-223.
- HOWELL, F. & R. ELLENDER (1984). Observations on growth and diet of *Argiope aurantia* Lucas (Araneidae) in a succession habitat. *Journal of Arachnology*, 12 : 29-36.
- INDYKIEWICZ, P., R. KUCHARSKI & P. ZALETA (1995). *Argiope bruennichi* Scopoli, 1772 (Aranei) on the eastern border of its distribution in Poland. *Przegląd Zoologiczny*, 39 : 87-89.
- JONSSON, L.J. & P. WILANDER (1999). Is the wasplike spider, *Argiope bruennichi*, established in Sweden? *Entomologisk Tidskrift*, 120 : 17-21.
- LINDEN van der, J. (2000). The expansion of the range of the spider *Argiope bruennichi* in The Netherlands (Araneae : Araneidae). *Nederlandse Faunistische Mededelingen*, 11 : 45-53.
- MALT, S., F.W. SANDER & G. SCHÄLLER (1990). Beitrag zur Nahrungsökologie ausgewählter Araneidae in Halbtrocken-

- rasen unter besonderer Berücksichtigung von *Argiope bruennichi* (Scop.). *Zoologische Jahrbücher. Zeitschrift für Systematic, Geographie und Biologie der Tiere*, 117 : 237-260.
- MCREYNOLDS, C.N. (2000). The impact of habitat features on web features and prey capture of *Argiope aurantia* (Araneae, Araneidae). *Journal of Arachnology*, 28 : 169-179.
- MOYES, N. (1997). *Argiope bruennichi*. First record for northern England. *Derbyshire Entomological Society Journal*, 127 : 2-5.
- NENTWIG, W. (1987). The prey of spiders. In : W. NENTWIG (ed.), *Ecophysiology of spiders*, Springer-Verlag, Berlin, Heidelberg, New York, London, Tokyo : 249-263.
- NYFFELER, M. & G. BENZ (1978). Die Beutespektren der Netzspinnen *Argiope bruennichi* (Scop.), *Araneus quadratus* Cl. und *Agelena labyrinthica* (Cl.) in Ödlandwiesen bei Zürich. *Revue Suisse de Zoologie*, 85 : 747-757.
- NYFFELER, M. & G. BENZ (1982). Eine Notiz zum Beutefangverhalten der Radnetzspinne *Argiope bruennichi* (Scopoli) (Araneae, Araneidae). *Revue Suisse de Zoologie*, 89 : 23-25.
- PASQUET, A. & R. LEBORGNE (1990). Prey capture efficiency and prey selection from insects intercepted by trap in four orb-weaving spider species. *Acta Oecologica*, 11 : 513-523.
- PASQUET, A. & R. LEBORGNE (1998). Behavioural tactic for prey capture and prey ingestion in two sympatric spiders. *Netherlands Journal of Zoology*, 48 : 39-52.
- PFLETSCHINGER, H. (1976). *Einheimische Spinnen. Die Webspinnen - Arten und Verhalten mit 120 Farbfotos*. Kosmos. Frankh Verlagshandlung, Stuttgart.
- POETZSCH, J. (1963). *Von der Brutfürsorge hemischer Spinnen*. Brehm-Bücherei 324. Zimsen Verlag. Wittenberg-Luthers-tadt.
- PUTS, C. (1989). Expansion territoriale de l'*Argiope fasciée* (*Argiope bruennichi* Scopoli) en Belgique et dans les régions voisines. *Verhandelingen van het Symposium "Invertebraten van België"*, Brussel 1988 : 193-198.
- RIECHERT, S.E. & J. LUCZAK, (1982). Spider foraging : behavioural responses to prey. In : P.N. WITT & J. ROVNER (eds.), *Biology of Spider Communication : Mechanisms and Ecological Significance*, Princeton Press, Princeton, NJ : 353-384.
- RIECHERT, S.E. & J.M. HARP (1987). Nutritional ecology of spiders. In : F.Jr. SLANSKY & J.G. RODRIGUEZ (eds.), *Nutritional ecology of insects, mites, spiders and related invertebrates*, John Wiley & Sons, New York, Chichester, Brisbane, Toronto, Singapore : 645-672.
- SCHARFF, N. & S. LANGEMARK (1997). *Argiope bruennichi* (Scopoli) in Denmark (Araneae; Araneidae). *Entomologiske Meddelelser*, 65 : 179-182.
- STARKS, P.T. (2002). The adaptive significance of stabilimenta in orb-webs : a hierarchical approach. *Annales Zoologici Fennici*, 39 : 307-315.
- URBAŃSKI, J. (1948). Distribution of *Argiope bruennichi* on Wolin Island and in the area of Poland. *Badania Fizjograficzne nad Polską Zachodnią*, 1 : 160-169.
- WEICKMANN, D. & A. GROBMAYER (1997). Der Einfluss des Winters auf die Population der Wespenspinne *Argiope bruennichi* Scopoli, 1772 (Araneae : Araneidae). *Arachnologisches Magazin*, 5 : 6-12.
- ZAR, J.H. (1999). *Biostatistical analysis*. 4<sup>th</sup> Edition. Prentice Hall, New Jersey.

Received: July 7, 2004

Accepted: December 10, 2004

